


THE SIGNS AND SYMPTOMS OF POMPE DISEASE

Pompe disease is a rare progressive disorder that is genetically inherited or passed on to children when both parents have a defective gene. People diagnosed with Pompe disease do not have enough of an enzyme called **acid alpha-glucosidase (GAA)**, or **acid maltase**. This enzyme is needed to break down **glycogen**, a form of sugar stored in muscle cells. When too much glycogen builds up in the muscle cells, the cells become damaged and the muscles cannot function properly. Because the enzyme is found in a part of the cell called the **lysosome**, Pompe disease is often referred to as a **lysosomal storage disorder**. Since the material stored is glycogen Pompe disease is also referred to as a glycogen storage disease (GSD), and since Pompe disease also affects the muscles, it is also called a **neuromuscular disorder**. Even though symptoms of Pompe disease may appear at any time from infancy through adulthood, patients receive the diagnoses of either infantile or late-onset Pompe disease. This brochure will describe the natural history of Pompe disease when untreated. Please note, there is a commercially available treatment accessible in most countries around the world that slows or stops the progression of Pompe in many patients. This will be discussed in other brochures.



Muscle cells (or fibers) are grouped in bundles. Each cell is made up of bands of **myofibrils**. Glycogen starts to build up in a part of the muscle cell called the **lysosome**. This causes the lysosomes to expand until they take up so much space that the muscle cell is damaged. Glycogen may also leak out of the cells and cause more damage to the muscle cells.

Frequently Asked Questions:

Q: How does the severity of Pompe disease relate to the age at which symptoms first appear?

A: All patients with Pompe disease share the same general disease course, namely the steady buildup of glycogen in muscle tissues leading to progressive muscle weakness. The broad spectrum of Pompe disease has the classic infantile onset at the severe end of the spectrum and the late onset at the other side of the spectrum. The severity of Pompe disease varies by age of onset, organ involvement and severity of muscular involvement (skeletal, respiratory, cardiac), and rate of progression.

In an attempt towards establishing a uniform terminology, Pompe disease is classified as follows:

- Classic Infantile Pompe disease
- Nonclassic Infantile Pompe Disease
- Late-Onset Pompe disease

Q: What are the signs and symptoms of classic and non-classic infantile Pompe disease?

A: Classic Infantile Pompe disease: This is the most aggressive and life-threatening form of the disease, and usually appears during the first 6 months of life. Extreme muscle weakness (myopathy) is the most noticeable sign. Babies with Pompe disease have poor muscle tone (hypotonia), look “floppy” and cannot hold up their heads. They typically are slow to acquire motor skills; they may lose previously acquired motor skills. They may never achieve the ability to sit up, crawl or stand. The muscle weakness progresses rapidly. Breathing, sucking, and swallowing become extremely difficult. The heart enlarges (cardiomegaly), the liver enlarges (hepatomegaly), and the tongue becomes enlarged (macroglossia). Affected infants may also fail to gain weight and fail to grow at the expected rate (failure to thrive) and have breathing problems. Increasing weakness of the heart muscle leads to heart and respiratory failure. If untreated, this form of Pompe disease leads to death in the first year of life. This represents the most severe form of the disease.

Mental development does not seem to be affected.

Nonclassic Infantile Pompe Disease: Usually appears by one year of age. It is characterized by delayed motor skills (such as rolling over and sitting up), and progressive muscle weakness. The heart may be abnormally large (cardiomegaly), which may result in heart failure similar to the most severely affected infants; the rate of progression may not be as rapid. Some patients have minimal to no cardiac involvement, with early muscle involvement as the primary presenting feature. The muscle weakness can lead to serious breathing problems, and if untreated, children with this form of Pompe disease will only live into early childhood.

Q: What are the signs and symptoms of late-onset Pompe disease?

A: Late-onset Pompe disease may not become apparent until later in childhood, adolescence, or adulthood. However, some present as early as the first year of life. Late-onset Pompe disease is usually milder than the infantile forms of this disorder and is less likely to involve the heart. Most individuals with late-onset Pompe disease experience progressive muscle weakness, especially in the legs and the trunk, including the muscles that control breathing.

THE SIGNS AND SYMPTOMS OF POMPE DISEASE

Those who first present symptoms of Pompe disease during childhood may have trouble keeping up with other children their age when they do physical activities. Adults may tire easily or get winded after exercising or climbing stairs. Some have low back pain. Enlargement of the heart or liver, a classic symptom of infantile onset Pompe disease, is rarely seen in late-onset Pompe disease. As muscle weakness increases, patients often start to use wheelchairs and may require assisted ventilation.

Pompe disease affects one of the most important muscles that we use for breathing - the diaphragm. The diaphragm is located just below the lungs and heart and separates the chest from the abdomen. As the diaphragm gets weaker, breathing becomes more difficult, especially while sleeping. Morning headaches and daytime sleepiness can result. In some cases, diaphragmatic weakness may be evident before any other significant weakness is noted. To learn more about assisted ventilation refer to the brochure titled Breathing Problems in Pompe Disease.

Another common identifying feature of Pompe disease is often a weakness of the legs or hips, which causes a swaying gait or waddle. People may have muscle aches and frequent falls. Babies may not learn to crawl, stand, walk, or meet other developmental milestones. As children get older they often develop a curvature of the spine: Lordosis (swayback), kyphosis (hunchback), or scoliosis (curved from side to side) that continues into adulthood.

It is important to remember that everyone has disease progression at a different rate and some children and adults have milder symptoms than others. Adult-onset Pompe disease can present as late as the second to sixth decade of life.

Q: Is fatigue an important feature of late-onset Pompe disease?

A: Fatigue is a frequently experienced symptom in adults with Pompe disease and may have a disabling impact on the lives of patients. Until recently, fatigue in Pompe disease did not receive much attention and was not structurally assessed. Fatigue is widespread among both mildly and severely affected adult patients with Pompe disease. The Fatigue Severity Scale (FSS) appears a useful tool in assessing fatigue in Pompe disease. The FSS is designed to differentiate fatigue from clinical depression, since both share some of the same symptoms. The FSS consists of answering a short questionnaire that requires the subject to rate his or her own level of fatigue.

Besides symptoms related to weakness of the skeletal and respiratory muscles, non-motor problems such as fatigue can also have a deep and disabling impact on the patients' lives. Fatigue is difficult to define, as it is often a non-specific and subjective complaint. Two suggested definitions are 'extreme and persistent tiredness, weakness or exhaustion mental, physical or both' and 'difficulty in initiation of or sustaining voluntary activities'. Although fatigue is a frequent symptom in many chronic disorders, it has received little attention in Pompe disease and was only sporadically reported.

For the best possible treatment of fatigue it is important to know why it is so notably present in Pompe disease. In a recent review the contribution of 'central' and 'peripheral' components to fatigue in neurological disorders has been discussed. In Pompe disease, a peripheral cause of fatigue, resulting from muscle weakness, is perhaps the most likely explanation. Especially relevant with respect to fatigue in Pompe disease is weakness of the respiratory muscles. Respiratory insufficiency may lead to fragmented sleep, which in turn may lead to daytime sleepiness and fatigue.

Fatigue is highly prevalent among adult patients with Pompe disease. It is present in both mildly and severely affected patients and is independent of disease duration. The FSS appears a useful tool for the assessment of fatigue in adult patients with Pompe disease. Further research is needed to unravel the pathophysiological mechanism and to identify targets for fatigue management.

THE SIGNS AND SYMPTOMS OF POMPE DISEASE

Q: Are pain and fatigue features of late-onset Pompe disease?

A: Although pain and fatigue are not specific for Pompe disease, they can have a strong impact on patients' lives. Data from the IPA- Erasmus MC Pompe Survey ^{Ref2} showed that 76% of the participating patients suffered from fatigue and that 46% experienced pain 'often' or 'always' in one or more areas of the body, mostly in the upper arms and legs.

Not much further research has been performed on pain in Pompe disease, while fatigue in adult patients was studied in more detail using the Fatigue Severity Scale (FSS). It turned out that fatigue was not only present among severely affected patients, but also among patients who were still mildly affected by the disease and had limited other complaints. It is important to note that fatigue is also an important first symptom among adult patients with Pompe disease.

To identify the best possible treatment of fatigue one first needs to know why it is such an important symptom in Pompe disease. In a review¹ it is discussed that fatigue in Pompe disease is most likely caused by muscle weakness, leading to increased muscle fatigability. However, patients do also experience a more generalized sense of fatigue, possibly initiated by feedback mechanisms from the brain to prevent excessive physical strain. As little is known about the cause of fatigue in Pompe disease, further research on this topic is needed.

Especially relevant with respect to fatigue in Pompe disease is weakness of the respiratory muscles. Respiratory insufficiency may lead to fragmented sleep, which in turn may lead to daytime sleepiness and fatigue. Pulmonary function testing in sitting and supine position is therefore indicated in patients with excessive fatigue, as well as a sleep study (polysomnography) to detect whether nightly hypoventilation or other sleep disorders could cause the present symptoms of fatigue.

Q: Why does it sometimes take so long to get the right diagnosis?

A: Pompe disease is very rare, affecting about 1 in 40,000 people. The infantile form of the disease may be easier to recognize because the unique and swiftly developing symptoms point toward a specific diagnosis.

Recognizing Pompe disease can be challenging as signs and symptoms may be diverse and shared with those of other disorders, such as Werdnig-Hoffmann disease, Polymyositis, Becker / Duchene muscular dystrophy, or limb-girdle muscular dystrophy. Diagnostic delays have been reported to average 7 years in older children and adults. A diagnosis of Pompe disease can be confirmed by measuring acid alpha-glucosidase (GAA) enzyme activity. It is now possible to accurately measure GAA activity in dried blood spots, mixed leukocytes, and lymphocytes.

Q: What health problems may occur with Pompe disease?

A: The progressive muscle weakness and breathing problems caused by Pompe disease can increase the risk for respiratory infections, sleep apnea (pauses in breathing while asleep), and difficulty in swallowing (dysphagia), as well as scoliosis, contractures (muscle tightness), and lower back pain. To learn more about the health problems caused by Pompe disease, refer the brochure titled Common Medical Concerns

Q: Is there a cure for Pompe disease?

A: Both first and Second Generation Enzyme replacement treatments are also now available in some countries worldwide.

THE SIGNS AND SYMPTOMS OF POMPE DISEASE

Researchers are also studying other ways to slow the progression of symptoms or cure the disease. For more on these approaches, see the Pompe connections brochure titled Medical Progress in Pompe Disease and Gene Therapy. Supportive therapies are also available to help manage the symptoms of the disease. Such therapies may bring comfort and relief and help people live their lives as fully as possible.

Ref 1

Pompe disease diagnosis and management guideline

[Genetics in Medicine](#)

ACMG Work Group on Management of Pompe Disease: [Priya S. Kishnani, MD,¹](#) [Robert D. Steiner, MD \(Chair\),²](#) [Deeksha Bali, PhD,¹](#) [Kenneth Berger, MD,³](#) [Barry J. Byrne, MD, PhD,⁴](#) [Laura Case, PT, DPT,¹](#) [John F. Crowley, JD, MBA,⁵](#) [Steven Downs, MD,⁶](#) [R. Rodney Howell, MD,⁷](#) [Richard M. Kravitz, MD,¹](#) [Joanne Mackey, CPNA,¹](#) [Deborah Marsden, MBBS,⁸](#) [Anna Maria Martins, MD,⁹](#) [David S. Millington, PhD,¹](#) [Marc Nicolino, MD, PhD,¹⁰](#) [Gwen O'Grady, MA,¹](#) [Marc C. Patterson, MD, FRACP,¹¹](#) [David M. Rapoport, MD,¹²](#) [Alfred Slonim, MD,¹³](#) [Carolyn T. Spencer, MD,⁴](#) [Cynthia J. Tiffit, MD, PhD,¹⁴](#) and [Michael S. Watson, PhD¹⁵](#)

Ref 2

Pompe disease in children and adults: natural course, disease severity and impact on daily life Results from an international patient survey
Marloes Hagemans

Where to Learn More: See Further information section

This publication is designed to provide general information in regard to the subject matter covered. It is distributed as a public service by the International Pompe Association, with the understanding that the International Pompe Association is not engaged in rendering medical or other professional services. Medicine is a constantly changing science. Human error and changes in practice make it impossible to certify the precise accuracy of such complex materials. Confirmation of this information from other sources, especially from one's physician is required.